Agriculture
Canada

Direction générale
de la recherche

Research
Branch

Bibliographie sur les altises phytophages des crucifères au Canada

A bibliography relevant to crucifer-feeding flea beetle pests in Canada

Bulletin technique nº 22
Technical Bulletin No. 22

Station de recherches
Saint-Jean-sur-Richelieu, Québec
Research Station

Canada

86-01614/

ISSN 0825-4559
ISBN 0-662-53962-1
No. de cat.: A-54-9/22

Bibliographie sur les altises phytophages des crucifères au Canada

A bibliography relevant to crucifer-feeding flea beetle pests in Canada

CHARLES VINCENT
Station de recherches
Research Station
Agriculture Canada
430 boul. Gouin Blvd.
Saint-Jean-sur-Richelieu
(Québec)
J3B 6Z8

LARRY BURGESS
Station de recherches
Research Station
Agriculture Canada
107 Science Crescent
Saskatoon
(Saskatchewan)
S7N 0X2

Direction générale de la recherche
Research Branch
Agriculture Canada
1985

On peut obtenir des exemplaires de cette publication à la
Copies of this publication are available from
Station de recherches
Research Station
Direction générale de la recherche
Research Branch
Agriculture Canada
C.P. 457
430 boul. Gouin Blvd.
Saint-Jean-sur-Richelieu (Québec)
J3B 6Z8

©Ministre des Approvisionnements et Services Canada 1985

©Minister of Supply and Services Canada 1985

Introduction

Les espèces Phyllotreta cruciferae (Goeze) et Phyllotreta striolata (F.) sont des ravageurs importants des crucifères cultivés dans diverses parties du monde. Cette bibliographie présente une liste des références que nous avons jugée en rapport avec la recherche sur ces insectes au Canada. Nous présentons également quelques références sur Psylliodes punctulata Melsh. Cette espèce est fréquemment présente sur nos crucifères cultivés, mais généralement à des niveaux sub-économiques. Quoique notre bibliographie soit incomplète, nous espérons que les références mentionnées seront utiles, en particulier celles publiées avant la venue des banques de données informatisées. Le lecteur désireux de lire davantage sur les problèmes reliés aux altises est prié de consulter la bibliographie de Jourdheuil (1963) sur les Chrysomelidae.

Remerciements

Nous remercions Ghyslaine Brodeur, de la Station de recherche de Saint-Jean-sur-Richelieu, pour sa patience lors du traitement de texte du manuscrit, et Diana Kichuck et Gail Charabin de la bibliothèque de la Station de recherche de Saskatoon pour leur précieuse assistance lors de la préparation de cette bibliographie.

Introduction

The flea beetles Phyllotreta cruciferae (Goeze) and Phyllotreta striolata (F.) are serious pests of cruciferous field and garden crops in various parts of the world. This bibliography lists references that we judge to be particularly relevant to research on these pests in Canada. Also included are some references to Psylliodes punctulata Melsh. This species is often present in cruciferous crops in this country, but usually only in subeconomic numbers. Our bibliography is undoubtedly incomplete, but we hope that the references listed will be useful, particularly those published prior to the advent of computer listing and searching. For additional references to other problems on flea beetles, the reader is referred to the bibliography on Chrysomelidae in Jourdheuil (1963).

Acknowledgements

We thank Ghyslaine Brodeur of the Saint-Jean-sur-Richelieu Research Station for patiently processing the manuscript, and Diana Kichuk and Gail Charabin of the library of the Saskatoon Research Station for their valuable assistance in preparing this bibliography.

References

Anonymous. 1891. Turnip flea-beetle (Phyllotreta vittata, Fab.). Can. Dep. Agric. Bull. No. 11, p. 30.

Anonymous. 1905. "Flea" beetles (Halticinae). Great Br. Minist. Agric. Fish. Food Leafl. No. 3, 4 p.

Anonymous. 1978. Flea beetles. Great Br. Minist. Agric. Fish. Food Advs. Leafl. No. 109, 4 p.

Anonymous. 1980. Oilseed rape. Great Br. Minist. Agric. Fish. Food Booklet No. 2278, 26 p.

Alfieri, A. 1976. The Coleoptera of Egypt. Mem. Soc. Entomol. Egypte 5: 1-362.

Allard, E. 1860. Essai monographique sur les Galérucites anisopodes (Latr.) ou description des altises d'Europe et des bords de la mer Méditerranée. Ann. Soc. Entomol. Fr. (série 3) 8: 370-418.

Allard, E. 1866. Monographie des Alticides, tribu de la famille des Phytophages. L'Abeille 3: 169-508.

Allen, A.A. 1976. Notes on some British Chrysomelidae (Col.) including amendments and additions to the list. Entomol. Rec. J. Var. 88: 220-25.

Allen, A.A. 1977. A recent occurrence of Phyllotreta vittata F. (sinuata Auct. Brit.) (Col.: Chrysomelidae). Entomol. Rec. J. Var. 89: 332.

Arnett, R.H. 1968. The beetles of United States (a manual of identification). American Entomological Institute, Ann Arbor, Mich., 1112 p.

Bakhetia, D.R.C. and K.S. Labana. 1978. Insect resistance in Brassica crops. Crop Improv. 5: 95-103.

Balsbaugh, E.U. Jr. and K.L. Hays. 1972. The leaf beetles of Alabama (Coleoptera: Chrysomelidae). Ala. Agric. Exp. Stn., Auburn, Bull. 441, 223 p.

Balsbaugh, E.U., R.W. Kieckhefer and E.L. Miller. 1967. Aerial collection of flea beetles (Chrysomelidae: Alticinae) in South Dakota. Proc. North Cent. Branch Entomol. Soc. Am. 22: 154-58.

Baly, J.S. 1874. Catalogue of the phytophagous Coleoptera of Japan with descriptions of species new to science. Trans. Entomol. Soc. Lond. Ser. 4: 161-217.

Baly, J.S. 1877a. Descriptions of new genera and of uncharacterized species of Halticinae. Trans. Entomol. Soc. Lond. 1877, Pt. II: 157-84.

Baly, J.S. 1877b. Descriptions of new genera and of uncharacterized species of Halticinae. Trans. Entomol. Soc. Lond. 1877, Pt. IV: 283-323.

Banerjee, S.N. and A. Basu. 1955. Phyllotreta chotanica Duvivier, a new pest of mustard in West Bengal. Curr. Sci. 24: 14-15.

Barber, H.S. 1947. Diabrotica and two new genera (Coleoptera: Chrysomelidae). Proc. Entomol. Soc. Wash. 49: 151-61.

Bardner, R. and W.E. Taylor. 1970. Effects of flea beetles on the yield of radish. Entomol. Exp. Appl. 13: 54-60.

Bargali, di P. 1878. La flora delle altiche in Europa. Bull. Soc. Entomol. Ital. 10: 43-73, 126-53, 204-13.

Batra, H.N. 1969. Food plants, bionomics and control of flea beetles. Indian Farming 19(3): 38-40.

Bedel, L. 1889-92. Faune des Coléoptères du bassin de la Seine. Tome V. Publ. Soc. Entomol. Fr., Paris, 423 p.

Beirne, B.P. 1971. Pest insects of annual crop plants in Canada. III. Coleoptera. Mem. Entomol. Soc. Can. No. 78: 71-124.

Bird, R.D. 1948. Flea beetles. Can. Dep. Agric. Processed Publ. No. 83. 3 p.

Blackwelder, R.E. 1944. Checklist of the coleopterous insects of Mexico, Central America, the West Indies, and South America, Part 4. Smithson. Inst. U.S. Nat. Mus. Bull. No. 185: 551-763.

Blatchley, W.S. 1910. An illustrated descriptive catalogue of the Coleoptera or beetles (exclusive of the Rhynchophora) known to occur in Indiana. The Nature Publishing Co., Indianapolis, 1386 p.

Bodor, J. and K. Balazs. 1970. Possibilities of the control of insect pests in Brassica cultures. Novenyved Korszerusitese 4: 79-104. (Hungarian; English summary).

Bondarenko, N. 1982. Prospects of biological control of pests in cruciferous vegetable crops in the USSR. Acta Entomol. Fenn. 40(0): 3-5.

Bonnemaison, L. 1965. Insect pests of crucifers and their control. Annu. Rev. Entomol. 10: 233-56.

Brett, C.H. and J.D. Rudder. 1966. Resistance of 30 commercial cruciferous varieties to the striped flea beetle, Phyllotreta striolata. J. Econ. Entomol. 59: 769.

Brett, C.H. and M.J. Sullivan. 1974. The use of resistant varieties and other cultural practices for control of insects on crucifers in North Carolina. N.C. Agric. Exp. Stn., Raleigh, Bull. 449, 30 p.

Brook, A.M. 1980. Flea beetles. Great Br. Minist. Agric. Fish. Food, Leafl. No. 109, 5 p.

Brown, W.J. 1967. Notes on the extralimital distribution of some species of Coleoptera. Can. Entomol. 99: 85-93.

Burgess, L. 1977a. Flea beetles (Coleoptera: Chrysomelidae) attacking rape crops in the Canadian Prairie provinces. Can. Entomol. 109: 21-32.

Burgess, L. 1977b. Geocoris bullatus, an occasional predator on flea beetles (Hemiptera: Lygaeidae). Can. Entomol. 109: 1519-20.

Burgess, L. 1980a. The horseradish flea beetle in Saskatchewan. Blue Jay. 38: 11-13.

Burgess, L. 1980b. Predation on adults of the flea beetle Phyllotreta cruciferae by lacewing larvae (Neuroptera: Chrysopidae). Can. Entomol. 112: 745-46.

Burgess, L. 1981a. An attractant trap for crucifer-feeding flea beetles. Canadex 620.

Burgess, L. 1981b. Crucifer-feeding flea beetles (Coleoptera: Chrysomelidae) occurring in the province of Saskatchewan, Canada. Coleopt. Bull. 35: 307-09.

Burgess, L. 1981c. Flea beetles that attack rape and cruciferous garden crops in the Prairie Provinces. Canadex 622, 625.

Burgess, L. 1981d. Occurrence of flea beetles outside the western rape growing area. Canadex 620.149.

Burgess, L. 1981e. Winter sampling to determine overwintering sites and estimate density of adult flea beetle pests of rape (Coleoptera: Chrysomelidae). Can. Entomol. 113: 441-47.

Burgess, L. 1982a. Occurrence of Phyllotreta striolata, the striped flea beetle, in open prairie, forest and parkland of Saskatchewan (Coleoptera: Chrysomelidae). Can. Entomol. 114: 439-46.

Burgess, L. 1982b. Occurrence of some flea beetle pests of parkland rapeseed crops in open prairie and forest in Saskatchewan (Coleoptera: Chrysomelidae). Can. Entomol. 114: 623-27.

Burgess, L. 1982c. Predation on adults of the flea beetle Phyllotreta cruciferae by the western damsel bug, Nabis alternatus (Hemiptera: Nabidae). Can. Entomol. 114: 763-64.

Burgess, L. 1983. How flea beetle pests of canola crops overwinter. Canadex 622.149.

Burgess, L. 1984. Changes in the relative abundance of the flea beetles Phyllotreta striolata and Phyllotreta cruciferae (Coleoptera: Chrysomelidae) in Saskatchewan parkland with increasing distance from the boreal forest. Can. Entomol. 116: 653-56.

Burgess, L. and D.T. Spurr. 1984. Distribution and abundance of overwintering flea beetles (Coleoptera: Chrysomelidae) in a grove of trees. Environ. Entomol. 13: 941-44.

Burgess, L. and J.E. Wiens. 1976. Maintaining a colony of the striped flea beetle, Phyllotreta striolata (Coleoptera: Chrysomelidae), in the greenhouse. Can. Entomol. 108: 53-55.

Burgess, L. and J.E. Wiens. 1980. Dispensing allyl isothiocyanate as an attractant for trapping crucifer-feeding flea beetles. Can. Entomol. 112: 93-97.

Chagnon, G. 1917. A preliminary list of the insects of Quebec, Part III - Coleoptera. Suppl. Rep. Que. Soc. Prot. Plants, 277 p.

Chagnon, G. and A. Robert. 1962. Principaux Coléoptères de la Province de Québec. 2nd ed. Les Presses de l'Université de Montréal, Montréal, 440 p.

Chang, S.L. and C.C. Pegn. 1971. An investigation of chemical control of some important insect pests on Cruciferae in Singapore. Plant Prot. Bull. Taipei 13: 110-20.

Chapuis, M.F. 1875. Histoire naturelle des insectes, genera des Coléoptères, famille des Phytophages, T. 11, Librairie Encyclopédique de Robert, Paris, 420 p.

Chittenden, F.H. 1895. The horse-radish flea beetle (Phyllotreta armoraciae Koch.). Insect Life 7: 404-06.

Chittenden, F.H. 1902. Some insects injurious to vegetable crops. U.S. Dep. Agric. Div. Entomol. Bull. 33, Washington, D.C., 117 p.

Chittenden, F.H. 1917. Horse-radish flea-beetle: its life history and distribution. U.S. Dep. Agric. Bull. 535, Washington, D.C., 16 p.

Chittenden, F.H. 1923. Notes on the distribution and habits of North American Phyllotreta (Coleop.). Proc. Entomol. Soc. Wash. 25: 131-39.

Chittenden, F.H. 1927. The species of Phyllotreta North of Mexico. Entomol. Am. (n.s.) 8: 1-63.

Chittenden, F.H. and H.O. Marsh. 1920. Western cabbage flea-beetle. U.S. Dep. Agric. Bull. No. 902, Washington, D.C., 21 p.

Cinitis, R. 1975. Ground beetles (Carabidae) in the agrocenoses of cruciferous field crops. Latv. Entomol. 17: 7-26. (In Latvian).

Cromartie, W.J. 1975. The effect of stand size and vegetational background on the colonization of cruciferous plants by herbivorous insects. J. Appl. Ecol. 12: 517-33.

Crotch, G.R. 1873. Materials for the study of the beetles of United States. Proc. Acad. Nat. Sci. Phila. 25: 19-83.

Crowson, R.A. 1981. The biology of the Coleoptera. Academic Press, Toronto, 802 p.

Curtis, J. 1883. Farm insects. John Van Voorst, Paternoster Row, London, 528 p.

Dejean, M. le Comte. 1837. Catalogue des Coléoptères de la collection de M. le Comte Dejean. 3rd ed. Méquignon-Marvis Père et Fils, Paris, 503 p.

Deletrez, H. and G. Roussel. 1979. Use of thiofanox (Dacamox 5G) in the control of rape flea-beetles. Meded. Fac. Landbouwwet. Riksuniv. Gent 44: 159-70. (In French; English summary).

Dixon, G.R., J.M. Hague and J. Braybrooks. 1975. Field reaction of swede cultivars to turnip crinkle virus. Plant Pathol. 24: 31-32.

Dobson, R.M. 1956. A note on the relative abundance of flea beetles (Phyllotreta Stephens and Psylliodes Berthold) on different cruciferous crops. J. Hortic. Sci. 31: 291-94.

Duckett, A.B. 1920. Annotated list of Halticini. Univ. Maryland Agric. Exp. Sta. Bull. No. 241: 111-55.

Dugas, A.L. 1938. The striped flea beetle. Louisiana State Univ. Bull. No. 298: 25-27.

DuPorte, E.M. 1914. The wavy striped flea beetle. Can. Entomol. 46: 433-35.

DuPorte, E.M. 1916. Insect notes, 1915. Annu. Rep. Que. Soc. Prot. Plants for 1915-16: 73-77.

Efremova, T.G. 1978. Pests of crucifers. Zashch. Rast. (Mosc.) 1978 (No. 2): 55-56. (In Russian).

El-Sawaf, S.K., S.M. Hammad and M.I. Abdel-Fattah. 1965. The morphology of the adult flea-beetle, Phyllotreta cruciferae Goeze (Coleoptera; Chrysomelidae), with a histological study of its alimentary canal. Alexandria J. Agric. Res. 13: 121-51.

Elsey, K.D. 1977. Parasitism of some economically important species of Chrysomelidae by nematodes of the genus Howardula. J. Invertebr. Pathol. 29: 384-85.

Fabricius, J.C. 1801. Systema Eleutheratorum, Tomus I, Kiliae, 506 p.

Feeny, P., K.L. Paauwe and N.J. Demong. 1970. Flea beetles and mustard oils: Host plant specificity of Phyllotreta cruciferae and P. striolata adults (Coleoptera: Chrysomelidae). Ann. Entomol. Soc. Am. 63: 832-41.

Finch, S. 1977. Monitoring insect pests of cruciferous crops. Proc. 1977 Bull. Crop Prot. Conf. - Pests and diseases, p. 219-26.

Findlay, G.M. 1978. Field evaluation of insecticides for controlling flea beetles in rapeseed. Annu. Conf. Manit. Agron. 1976: 45-47.

Fletcher, J. 1905. Insects injurious to grain and fodder crops, root crops and vegetables. Can. Dep. Agric. Bull. No. 52, 48 p.

Foudras, C. 1860. Histoire naturelle des Coléoptères de France. XI. Altisides. Magnin, Blanchard et Cie, Paris, 384 p.

Fougeroux, A. 1983. La lutte contre les ravageurs du sol en grandes cultures. Phytoma Déf. Cult. 346: 56-57.

Frost, S.W. 1949. Flea beetles attacking Cruciferae. J. Econ. Entomol. 42: 144-45.

Furth, D.G. 1976. Alticinae of Israel, with special reference to the systematic, zoogeographic, and host plant relationships of Phyllotreta and Longitarsus (Coleoptera: Chrysomelidae). Ph.D. Thesis, Cornell Univ., 363 p.

Furth, D.G. 1979. Zoogeography and host plant ecology of the Alticinae of Israel, especially Phyllotreta; with descriptions of three new species (Coleoptera: Chrysomelidae). Isr. J. Zool. 28: 1-37.

Furth, D.G. 1980. Inter-generic differences in the metafemoral apodeme of flea beetles (Chrysomelidae: Alticinae). Syst. Entomol. 5: 263-71.

Furth, D.G. 1982. The metefemoral spring of flea beetles (Chrysomelidae: Alticinae). Spixiana Suppl. (Muench.) 7: 11-27.

Garman, H. 1892. Cabbage flea-beetle. Annu. Rep. Kentucky Agric. Exp. Stn., 2, 1889: 23-25.

Garcia-Tejero, F.D. 1946. Las Pulguillas de las cruciferas cultivadas. Bol. Patol. Veg. Entomol. Agric. 14: 353-68.

Garcia-Tejero, F.D. 1955. Halticinos de interes agricola. Bol. Patol. Veg. Entomol. Agric. 21: 343-93.

Gerber, G.H. and C.E. Osgood. 1975. Collops vittatus (Coleoptera: Melyridae): a predator of flea beetle adults in rapeseed. Manit. Entomol. 9: 61-63.

Gibson, A. 1912. Reports on insects for the year. Annu. Rep. 49th Annual Meeting Entomol. Soc. Ont., p. 11-20.

Gibson, A. 1913. Flea-beetles and their control. Can. Dep. Agric. Entomol. Div. Circ. No. 2, 11 p.

Gibson, A. 1913-1914. The injurious flea-beetles of the Prov. of Quebec. Sixth Annu. Rep. Que. Soc. Prot. Plants from Insects and Fungus Diseases, 1913-14, p. 25-30.

Gibson, A. 1917. Cabbage insects. Ninth Annu. Rep. Que. Soc. Prot. Plants from Insects and Fungus Diseases, 1916-17, p. 30-41.

Gibson, A. 1934. Insects of the flower garden and their control. Can. Dep. Agric. Bull. 99 (n.s.), 69 p.

Glendenning, R. 1927. The cabbage flea-beetle and its control in British Columbia. Can. Dep. Agric. Pam. 80 (n.s.): 1-10.

Glick, P.A. 1939. The distribution of insects, spiders, and mites in the air. U.S. Dep. Agric. Tech. Bull. No. 673, 151 p.

Goble, H.W. 1965. Insects attacking agricultural crops and ornamental plants in Ontario in 1965. Proc. Entomol. Soc. Ont. 96: 5-6.

Goeze, J.A.E. 1777. Entomologische Beiträge zu des Ritter Linné zwölften Ausgabe des Natursystems. Weidmanns Erben und Reich, Leipzig, 348 p.

Görnitz, K. 1953. Untersuchungen über in Cruciferen enthaltene Insekten-Attraktivstoffe. Nachrichtenbl. Dtsh. Pflanzenschutzdienst (Berl.) 7: 81-95.

Goury, G. and J. Guignon. 1905-1907. Les insectes parasites des crucifères. Feuill. Jeunes Natl. 36: 97-99, 113-17, 125-32, 144-48; 37: 96-98, 209-13.

Gruev, B. 1977. Chrysomelidae (Coleoptera) collected in China by P.M. Hammond and in Korea by M. Josifov. Entomol. Rev. Jpn. 30: 19-24.

Gruev, B. 1978. A contribution to the knowledge of the Korean fauna of Chrysomelidae (Coleoptera). Entomol. Rev. Jpn. 32: 49-59.

Gupta, B.M. and J.C. Sharma. 1971. Efficacy of certain insecticidal control schedules against insect pests of cabbage. Madras Agric. J. 58: 889-91.

Haddock, M.J. 1942. Observations on the species of flea beetles infesting Brassica crops in the West of England. Annu. Rep. Long Ashton Agric. Hortic. Res. Stn. Univ. Bristol, p. 166-69.

Harris, C.R. 1973. Laboratory evaluation of candidate materials as potential soil insecticides. IV. J. Econ. Entomol. 66: 216-21.

Harris, C.R. and F. Gore. 1970. Laboratory tests on the contact toxicity of some insecticides to adults of the cabbage flea beetle, Phyllotreta cruciferae. J. Econ. Entomol. 63: 1025-26.

Harris, C.R. and G.B. Kinoshita. 1977. Influence of post-treatment temperature on the toxicity of pyrethroid insecticides. J. Econ. Entomol. 70: 215-18.

Harris, T.W. 1852. A treatise on some of the insects of New England which are injurious to vegetation. 2nd ed. White and Potter, Boston, 513 p.

Harukawa, C. and Y. Nisikawa. 1941. The resistance of the adult of Phyllotreta vittata to low temperatures in winter. Kontyu (Tokyo) 14: 200-07. (In Japanese).

Harukawa, C. and M. Tokunaga. 1938. Studies on the life history and bionomics of Phyllotreta vittata Fabricius I. Life history of Phyllotreta vitatta Fabricius. Mem. Coll. Agric. Kyoto Imp. Univ. No. 44 (Entomol. Ser. 5): 1-48.

Harukawa, C. and M. Tokunaga. 1939. Studies on the life history and bionomics of Phyllotreta vittata Fabricius II. On the hibernation of the flea-beetle. Kontyu (Tokyo) 13: 208-21. (In Japanese; English summary).

Hatch, M.H. 1971. The beetles of the Pacific Northwest. Univ. Wash. Publ. Biol. 16, Part V. Univ. Wash. Press. Seatle and London, 662 p.

Heikertinger, F. 1912. Myth of the brassica flea beetle: A word in justification of Haltica oleracea L. Zool. Bot. Ges. Wien 62: 69-81. (In German).

Heikertinger, F. 1913. Native Brassica flea beetle; a critical description of Haltincinae species which damage cultivated crucifers in Central Europe. Centralbl. Bakteriol. Parasitenkd. Infektionskr. 36: 98-127. (In German).

Heikertinger, F. 1918. Leaf mines of the crucifer pest Phyllotreta nemorum L. Kolon. Rundsch. 7: 13-15. (In German).

Heikertinger, F. 1925. Results of fifteen years of investigations on host plants of indigenous Halticinae. Entomol. Bl. 21: 119-31. (In German).

Heikertinger, F. 1954. Halticinae, Erdflohe, Flohkafer. Handb. Pflanzenkr. 5: 313-43.

Heikertinger, F. and E. Csiki. 1940. Chrysomelidae: Halticinae I. Partes 166 et 169, in S. Schenkling (ed.), Coleopterorum Catalogus. Dr. W. Junk, Berlin, 336 and 635 p.

Hicks, K.L. 1972. Host plant and habitat selection by crucifer flea beetles (Coleoptera: Chrysomelidae). Ph.D. Thesis, Cornell Univ., 151 p.

Hicks, K.L. 1974. Mustard oil glucosides: Feeding stimulants for adult cabbage flea beetles, Phyllotreta cruciferae (Coleoptera: Chrysomelidae). Ann. Entomol. Soc. Am. 67: 261-64.

Hicks, K.L. and J.O. Tahvanainen. 1974. Niche differentiation by crucifer-feeding flea beetles (Coleoptera: Chrysomelidae). Am. Midl. Nat. 91: 406-23.

Horn, G.H. 1889. A synopsis of the Halticini of boreal America. Trans. Am. Entomol. Soc. 16: 163-320.

Hossfeld, R. 1963. Synecological comparison of the fauna of winter and summer rape. Z. Angew. Entomol. 52: 209-54. (In German; English summary).

Howard, L.O. 1898. Injury by the western flea-beetle, Phyllotreta pusilla Horn. U.S. Dep. Agric. Div. Entomol. Bull. 10: 92-93.

Issi, I.V. and D.F. Radishcheva. 1979. Microsporidia-caused diseases of insect pests injuring cruciferous crops in Leningrad region. Zashch. Rast. No. 46: 19-23. (In Russian; English summary).

Jameson, H.R. 1950. Control of turnip flea beetle by benzene hexachloride drilled with the seed. Nature 165: 980.

Jameson, H.R. 1958. The mechanism of the control of turnip flea beetle by benzene hexachloride dressings on Brassicae seed. J. Sci. Food Agric. 9: 590-96.

Jameson, H.R. 1960. Gamma BHC liquid seed dressing for the control of turnip flea beetle. J. Sci. Food Agric. 11: 528-34.

Jeffery, H. and G.N. Foster. 1980. Control of turnip flea beetle damage to Brassica seedings, 1979. Ann. Appl. Biol. 94(1, suppl.): 4-5.

Jones, F.G.W. and M.G. Jones. 1964. Pests of field crops. Edward Arnold Ltd, London, p. 129-35.

Jourdheuil, P. 1957. Description d'un Ichneumonide (Mesechorini) parasite secondaire de diverses espèces de Phyllotreta (Hym. Ichneumonidae). Bull. Soc. Entomol. Fr. 62: 41-45.

Jourdheuil, P. 1960a. Influence de quelques facteurs écologiques sur les fluctuations de population d'une biocénose parasitaire. Ann. Epiphyt. 11: 445-658.

Jourdheuil, P. 1960b. Remarques sur le nombre de générations de quelques Phyllotreta (Col. Chrysomelidae). Bull. Soc. Entomol. Fr. 65: 126-31.

Jourdheuil, P. 1963. Sous famille des Halticinae, p. 762-854. In A.S. Balachowsky (ed.), Entomologie Appliquée à l'Agriculture. Tome 1, Vol. 2, Coléoptères. Masson et Cie, Paris, 1391 p.

Kareiva, P. 1982a. Exclusion experiments and competitive release of insects feeding on collards. Ecology 63: 696-704.

Kareiva, P. 1982b. Experimental and mathematical analyses of herbivore movement: Quantifying the influence of plant spacing and quality on foraging discrimination. Ecol. Monogr. 52: 261-82.

Kareiva, P. 1983. Local movement in herbivorous insects: Applying a passive diffusion model to mark recapture field experiments. Oecologia 57: 322-27.

Karavyanskii, N.S. and N.M. Chyornykh. 1972. Chemical and agronomic measures for the control of crucifer flea beetles in common turnips. Dokl. Vses. Akad. S.-Kh. Nauk Im. V.I. 6: 21-22. (In Russian).

Kinoshita, G.B. 1976. Biology and control of Phyllotreta cruciferae (Goeze) (Coleoptera: Chrysomelidae) in Southwestern Ontario. Ph.D. Thesis, Dep. of Environmental Biology, Univ. of Guelph. 107 p.

Kinoshita, G.B., C.R. Harris, H.J. Svec and F.L. McEwen. 1978. Laboratory and field studies on the chemical control of the crucifer flea beetle, Phyllotreta cruciferae (Coleoptera: Chrysomelidae), on cruciferous crops in Ontario. Can. Entomol. 110: 795-803.

Kinoshita, G.B., H.J. Svec, C.R. Harris and F.L. McEwen. 1979. Biology of the crucifer flea beetle, Phyllotreta cruciferae (Coleoptera: Chrysomelidae), in Southwestern Ontario. Can. Entomol. 111: 1395-407.

Koshihara, T. 1981. Insecticides, vegetables and ornamental crops. Jpn. Pestic. Inf. 38: 27-29.

Kostromitin, V.B. 1978. Damage caused by cruciferous-feeding flea beetles. Zashch. Rast. 1978(No. 7): 36. (In Russian).

Kral, J. 1973. Results of the zoological researches of Dr. Z. Kaszab in Mongolia. 329. Alticidae. III. Coleoptera. Folia Entomol. Hung. 26(n.s.): 111-34. (In German; English summary).

Kroeker, G. 1977. Seed dressing of Brassica oilseeds, experiences from 1976. Plant Prot. Conf. 1977 Uppsala: 23-25. (In Swedish).

Lamb, R.J. 1980. Hairs protect pods of mustard (Brassica hirta 'Gisilba') from flea beetle feeding damage. Can. J. Plant Sci. 60: 1439-40.

Lamb, R.J. 1983. Phenology of flea beetle (Coleoptera: Chrysomelidae) flight in relation to their invasion of canola fields in Manitoba. Can. Entomol. 115: 1493-502.

Lamb, R.J. 1984. Effects of flea beetles, Phyllotreta spp. (Chrysomelidae: Coleoptera), on the survival, growth, seed yield and quality of canola, rape and yellow mustard. Can. Entomol. 116: 269-80.

Lamb, R.J. and W.J. Turnock. 1982. Economics of insecticidal control of flea beetles (Coleoptera: Chrysomelidae) attacking rape in Canada. Can. Entomol. 114: 827-40.

Larsen, L.M., J.K. Nielsen and H. Sorensen. 1982. Identification of 30-2-0 beta-D xylo pyranosyl-beta-D-galactopyranosyl flavenoids in horseradish leaves acting as feeding stimulants for a flea betle. Phytochemistry (Oxf.) 21: 1029-33.

Latheef, M.A. and J.A. Ortiz. 1984. Influence of companion herbs on Phyllotreta cruciferae (Coleoptera: Chrysomelidae) on collard plants. J. Econ. Entomol. 77: 80-82.

Latheef, M.A., J.H. Ortiz and A.Q. Sheikh. 1984. Influence of intercropping on Phyllotreta cruciferae (Coeloptera: Chrysomelidae) populations on collard plants. J. Econ. Entomol. 77: 1180-84.

Latreille, P.A. 1812. Histoire naturelle, générale et particulière des crustacés et des insectes. Tome 12. F. Dufort, Paris, 424 p.

Laumond, C., H. Mauléon and A. Kermarrec. 1979. Données nouvelles sur le spectre d'hôtes et le parasitisme du nématode entomophage Neoaplectana carpocarpsae. Entomophaga 24: 13-27.

Lazorko, W. 1973. Three species of Coleoptera new to British Columbia. J. Entomol. Soc. B.C. 70: 41.

Lechapt, G. 1980. Methods of forecasting insect infestation in oilseed rape. Bull. OEPP (Org. Eur. Medit. Prot. Plant.) 10: 119-28. (In French; English , Russian summaries).

Lee, Y.W. and N.D. Westcott. 1980. Direct analysis of carbofuran and 3 hydroxy carbofuran in rape plants, Brassica napus and Brassica compestris, by reverse phase high pressure liquid chromatography. J. Agric. Food Chem. 28: 719-22.

Leng, C.W. 1920. Catalogue of the Coleoptera of America, north of Mexico. John D. Sherman, Jr., Mount Vernon, New York, 470 p.

Leonardi, C. 1975. Bio geographic data on Alticinae of Sardinia with preliminary descriptions of a new Aphthona species (Coleoptera Chrysomelidae). Atti. Soc. Ital. Sci. Nat. Mus. Civ. Stor. Nat. Milano 116(1-2): 3-14. (In Italian; English summary).

Liashenko, L.I. 1980. Turnip flea beetles. Zashch. Rast. 1980 (No. 10): 53.

Loan, C.C. 1967. Studies on the taxonomy and biology of the Euphorinae (Hymenoptera: Braconidae). II. Host relations of six Microctonus species. Ann. Entomol. Soc. Am. 60: 236-40.

Lopatin, I.K. 1960. Materials on the fauna and ecology of leaf-beetles (Coleoptera, Chrysomelidae) of the southern bank of the Dnieper River. Entomol. Obozr. 39: 629-42. (In Russian; English summary).

Lundberg, S. 1971. Further finds of some interesting beetle species (Thinobiuos-brevipennis Kies, Atheta-trybomi J. Sahlb., Atheta-wireni Brd., and Phyllotreta-zimmermanni Crotch). Entomol. Tidskr. 92: 267-70. (In Swedish).

Matsumoto, Y. 1970. Volatile organic sulphur compounds as insect attractants with special reference to host selection. p. 133-160 in D.L. Wood, R.M. Silverstein and M. Nakajima (eds.), Control of insect behavior by natural products. Academic Press, New York, 345 p.

McMahon, E. 1942. Observations on flea beetles which attack cruciferous crops. J. Dep. Agric. (Repub. Irel.) 39: 80-83.

Medvedev, L.N. 1979. Further contribution to the knowledge of Chrysomelidae from Mongolia (Coleoptera). Folia Entomol. Hung. 32(2): 161-67.

Meisner, J. and B.K. Mitchell. 1983. Phagodeterrency induced by two cruciferous plants in adults of the flea beetle Phyllotreta striolata (Coleoptera: Chrysomelidae). Can. Entomol. 115: 1209-14.

Mekhonoshina, Z.G. 1976. Rogor against crucifer flea-beetles. (Abstract) Zashch. Rast. 1976 (No. 5): 51. (In Russian).

Melsheimer, F.E. 1847. Description of new species of Coleoptera of the United States. Proc. Acad. Nat. Sci Phila. 3: 158-81.

Miles, H.W. 1935. The control of flea beetles with a naphthalene-silica dust. J. Minist. Agric. (Great Br.) 41: 1079-83.

Miles, H.W. 1943. Turnip flea beetles. Agriculture, Great Br. Minist. Agric., Fish and Food 50: 26-38.

Miles, H.W. 1944. Experiments on the control of flea beetles on Brassicae in the Western Province. Annu. Rep. Long Ashton Res. Stn. Univ. Bristol, p. 145-49.

Milliron, H.E. 1953. A European flea beetle injuring crucifers in North America. J. Econ. Entomol. 46: 179.

Milliron, H.E. 1958a. Economic insects and allied pests of Delaware. Univ. Del. Agric. Exp. Stn. Bull. No. 321: 26-28, 51, 74-75, 81.

Milliron, H.E. 1958b. Introduced flea beetle, Phyllotreta cruciferae Goeze. Del. Agric. Exp. Stn. Bull. 321: 26, 74-75, 81-82.

Minoranskiy, V.A. 1973. Leaf-beetles (Coleoptera: Chrysomelidae) damaging beet in the North Caucasus. Entomol. Rev. 52: 347-56.

Moreton, B.D. 1945. On the migration of flea beetles (Phyllotreta spp.) (Col., Chrysomelidae) attacking Brassica crops. Entomol. Mon. Mag. 81: 59-60.

Nakat, S.S. and K.M. Peswani. 1972. Relative efficacy of some granular systemic insecticides against important pests of mustard. Indian J. Entomol. 34: 118-22.

Narayanan, E.S., B.R. Subba Rao, M. Ramachandra Rao and A.K. Sharma. 1960. Biology and morphology of the immature stages of Microctonus indicus, new species (Braconidae: Hymenoptera) a parasite of Phyllotreta cruciferae Goeze (Chrysomelidae: Coleoptera). Proc. Indian Acad. Sci. Sect. B. 51: 280-87.

Newton, H.C.F. 1928a. The biology of flea-beetles (Phyllotreta) attacking cultivated cruciferae. J. S.-East. Agric. Coll. Wye, Kent, Great Br., No. 25: 90-115.

Newton, H.C.F. 1928b. Experiments on the control of flea-beetles of the genus Phyllotreta injurious to cultivated crucifer crops. J. S.-Est. Agric. Coll. Wye, Kent, Great Br., No. 25: 116-40.

Newton, H.C.F. 1929. Observations on the biology of some flea beetles of economic importance. J. S.-East Agric. Coll. Wye, Kent. Great Br. No. 26: 145-64.

Newton, H.C.F. 1931. Notes on some parasites reared from flea-beetles of the genus Phyllotreta (Chrysomelidae). Entomol. Mon. Mag. 66: 82-84.

Newton, H.C.F. 1934. Insect pests at Rothamsted and Woburn. 1932-3. Rep. Rothamsted Exp. Stn. 1933: 51-53.

Newton, H.C.F. 1935. Insect pests at Rothamsted and Woburn. 1933-4. Rep. Rothamsted Exp. Stn. 1934: 71-73.

Nielsen, J.K. 1977. Host plant relationships of Phyllotreta nemorum L. (Coleoptera: Chrysomelidae). Z. Angew. Entomol. 84: 396-407.

Nielsen, J.K. 1978a. Host plant discrimination within Cruciferae: feeding responses of four leaf beetles (Coleoptera: Chrysomelidae) to glucosinolates, cucurbitacins, and cardenolides. Entomol. Exp. Appl. 24: 41-54.

Nielsen, J.K. 1978b. Host plant selection of monophagous and oligophagous flea beetles feeding on crucifers. Entomol. Exp. Appl. 24: 562-69.

Nielsen, J.K., L. Dalgaard, L.M.. Larsen and H. Sorensen. 1979. Host plant selection of the horse-radish flea beetle Phyllotreta armoraciae (Coleoptera: Chrysomelidae): feeding responses to glucosinolates from several crucifers. Entomol. Exp. Appl. 25: 227-39.

Nielsen, J.K., L.M. Larsen and H. Sorensen. 1977. Cucurbitacin E and I in Iberis amara; feeding inhibitors for Phyllotreta nemorum. Phytochemistry 16: 1519-22.

Nielsen, J.K., L.M.. Larsen and H. Sorensen. 1979. Host plant selection of the horseradish flea beetle Phyllotreta armoraciae (Coleoptera: Chrysomelidae): identification of two flavonol glycosides stimulating feeding in combination with glucosinolates. Entomol. Exp. Appl. 26: 40-48.

Nonveiller, G. 1960. Flea beetles harmful to useful plants in Serbia, species verified and their plant hosts. Belgrade. Inst. Za Zastitu Bilja. Posebna Izd. 10, 55 p. (In Serbo-Croatian; French summary).

Oatman, E.R. and G.R. Platner. 1969. An ecological study of insect populations on cabbage in Southern California. Hilgardia 40: 1-40.

Oldham, J.N. 1933. On Howardula phyllotretae n.sp., a nematode parasite of flea beetles (Chrysomelidae; Coleoptera), with some observations on its incidence. J. Helminthol. 11: 119-36.

Oldham, J.N. 1935. Further observations on the incidence of parasitism of flea beetles by the nematode, Howardula phyllotretae. J. Helminthol. 13: 163-66.

Osgood, C.E. 1974. Flea beetles on rape. Canadex 149.622.

Osgood, C.E. 1975a. Aspects of pest management for flea beetles in rape. Pap. presented at Manit. Agron. Annu. Conf. Dec. 1975: 51-54.

Osgood, C.E. 1975b. Damage assessment as part of flea beetle management on rape. Proc. 22nd Annu. Meet. Agric. Pestic. Soc. (Canada).

Parker, W.B. 1910. Some insects injurious to truck crops. The life history and control of the hop flea-beetle (Psylliodes punctulata Melsh.). Bull. U.S. Dep. Agric. Bur. Entomol. 82: 33-58.

Patil, S.P. and R.N. Pokharkar. 1979. Some new records of insect pests infesting cruciferous vegetable crops in Maharashtra State, India. J. Maharashtra Agric. Univ. 4: 222-23.

Peregudt, M.F. 1977. An experiment in cultivating Crambe steveniana as a new vegetable crop. Rastit. Resur. 13: 94-99. (In Russian).

Petersen, B. 1977. Pollination of Thlaspi alpestre by selfing and by insects in the alpine zone of Colorado. Arct. Alp. Res. 9: 211-15.

Pimentel, D. 1961. Competition and the species per-genus structure of communities. Ann. Entomol. Soc. Am. 54: 323-33.

Pjanic, E. and K. Thaler. 1981. Leaf beetles (Chrysomelidae: Halticinae) in montane field habitats near Innsbruck, Austria. Ber. Naturwiss. -Med. Ver. Innsb. 68: 137-44. (In German).

Pjatakova, V.L. 1928. Flea beetles. Bull. Mleev Hortic. Exp. Sta. No. 11: 1-75. (In Russian; French summary).

Proeseler, G. and K. Schmellzer. 1977. Laboratory and field tests on the transmission of Erysimum latent virus by Phyllotreta spp. Zentalbl. Bakteriol. Parasitenkd. Infektionskr. Hyg. Zweite 132: 716-21. (In German).

Provancher, L. 1877. Petite faune entomologique du Canada. Vol. I - Les Coléoptères. Presses de C. Darveau, Québec, 785 p.

Putnam, L.G. 1969. Rapeseed's "Big 3" insect pests. Can. Agric. 14(3): 22-23.

Putnam, L.G. 1977. Response of four Brassica seed crop species to attack by the crucifer flea beetle, Phyllotreta cruciferae. Can. J. Plant Sci. 57: 987- 89.

Putnam, L.G. and L. Burgess. 1977. Insect pests of rape and mustard. 1977. In Rapeseed Assoc. Can. Publ. No. 48, p. 3-20.

Reed, H.E. and R.A. Byers. 1981. Flea beetles attacking forage kale, Brassica oleracea). Effect of carbofuran and tillage methods. J. Econ. Entomol. 74: 334-37.

Riley, E.G. and W.R. Enns. 1979. An annotated checklist of Missouri leaf beetles (Coleoptera: Chrysomelidae). Mo. Acad. Sci. Trans. 13: 53-83.

Roberts, J.E. Sr. and M.A. Saluta. 1984. Cabbage flea beetle, Phyllotreta cruciferae (Coleoptera: Chrysomelidae), cabbage looper, Trichoplusia ni (Lepidoptera: Noctuidae) and imported cabbage worm, Pieris rapae (Lepidoptera: Pieridae), control on cabbage, Brassica oleracea capitata, with at-planting carbofuran treatments. Va. J. Sci. 35: 69.

Root, R.B. 1973. Organization of a plant-arthropod association in simple and diverse habitats: the fauna of collards (Brassica oleracea). Ecol. Monogr. 43: 95-124.

Root, R.B. 1975. Some consequences of ecosystem texture. In S.A. Levin (Ed.). Ecosystem analysis and prediction. Soc. Ind. Appl. Math., Philadelphia, 337 p.

Root, R.B. and J.J. Skelsey. 1969. Biotic factors involved in crucifer aphid outbreaks following insecticide application. J. Econ. Entomol. 62: 223-33.

Root, R.B. and J.O. Tahvanainen. 1969. Role of winter cress, Barbarea vulgaris, as a temporal host in the seasonal development of the crucifer fauna. Ann. Entomol. Soc. Am. 62: 852-55.

Ruter, G. 1978. Note on Phyllotreta armoraciae Koch (Coleoptera: Chrysomelidae). Entomologiste (Paris) 34: 150. (In French).

Rymer-Roberts, A.W. 1918. Turnip flea-beetles - wireworms. Entomologist (Lond.) 1918: 187.

Scherer, G. von. 1969. Die Alticinae des Indischen Subkontinentes (Coleoptera-Chrysomelidae). Pac. Insects Monogr. 22: 1-251.

Scherer, G. 1982. First International Alticinae Symposium, Munich, West Germany. Spixiana (Muench.) 7: 7-10. (In English).

Selivanon, S.N. 1935. Study of the ecology of Phyllotreta cruciferae Goeze in the zone of its injurious activity. Plant Prot. Leningrad, 1935: 96-98. (In Russian).

Shimer, H. 1869. The wavy-striped flea-beetle. Am. Nat. 2: 514-17.

Shukla, D.D., G. Proeseler and K. Schmelzer. 1975. Studies on viruses and virus diseases of cruciferous plants. XVIII. Beetle transmission and some new natural hosts of Erysimum latent virus. Acta. Phytopathol. Acad. Sci. Hung. 10: 211-15.

Smith, E.H. 1973. Systematic revision of the maculate species of the genus Phyllotreta (Chev.) of America North of Mexico (Coleoptera: Chrysomelidae, Alticinae). Ph.D. Thesis, Dep. of Entomology, The Ohio State Univ., 221 p.

Smith, E.H. 1977. Clarification notes on two Fabrician chrysomelid type specimens (Coleoptera: Chrysomelidae). Coleopt. Bull. 31: 87-90.

Smith, E.H. 1979. Genus Tanygoster Blatchley, a new synonymn of Phyllotreta Chevrolat (Coleoptera: Chrysomelidae: Alticinae). Coleopt. Bull. 33: 359-62.

Smith, K.M. and R. Markham. 1946. An insect vector of the turnip yellow mosaic virus. Nature 158: 417.

Smith, O.J. and A. Peterson. 1950. Microctonus vittatae, a parasite of adult flea beetles, and observations on hosts. J. Econ. Entomol. 43: 581-85.

Snelling, E.T. 1867. Note. Pract. Entomol. 2: 119.

Sommer, G. 1981. Biologie und Parasitenkomplex der Halticinen Gattung Phyllotreta. Doctoral Thesis. Albert-Ludwigs-Univ. zu Freiburg in Breisgau, 317 p.

Sommer, G. 1981. Biology and parasites of Phyllotreta spp. (Coleoptera: Halticinae). Commonw. Inst. Biol. Control. Eur. Stn., Delemont, Switz. Rep. 45 p.

Srivasatava, O.S. 1974. A new host record for Phyllotreta cruciferae Goeze (Coleoptera: Chrysomelidae). Indian J. Hortic. 31: 121.

Srivastava, A.S., B.K. Awasthi and Y.P. Singh. 1973. Incidence of flea beetle, Phyllotreta cruciferae Goez., (Coleoptera, Chrysomelidae) on radish crop. Indian J. Zootomy 14: 67-68.

Stear, J.R. 1919. Flea beetles as pests of the garden. (Abstr.) Rev. Appl. Entomol. (A) 7: 147-48.

Steinhausen, W.R. 1982. Status of Palearctic Alticinae larvae research. Spixiana Suppl. (Muench.) (7): 39-42.

Stenmark, A. 1979. Research concerning the control of flea beetles in spring rape. Vaxtskyddsnotiser-sver. Lantbruksuniv. 43: 48-50. (In Swedish; English summary).

Stewart, R.P. 1943. A modification of the flea beetle trap. J. Roy. Hortic. Soc. (Lond.) 68: 275.

Strickland, E.H. and B. Hocking. 1950. Insects of the Alberta farmstead. Univ. Alberta Agric. Ext. Bull. 55, 118 p.

Suzuki K. 1974. Ovariole number in the family Chrysomelidae (Insecta: Coleoptera). J. Coll. Liberal Arts (Toyama Univ., Jpn.) 7: 53-70.

Suzuki, K. 1978. Historical review of studies of the internal reproductive system in the family Chrysomelidae s.L. (Coleoptera). J. Coll. Liberal Arts (Toyama Univ., Jpn.) 11: 61-80.

Suzuki, K. and A. Hara. 1976. Comparative study of the egg size in relation to the egg number in the family Chrysomelidae (Insecta: Coleoptera). J. Coll. Liberal Arts (Toyama Univ., Jpn.) 9: 39-81.

Tahvanainen, J.O. 1971. Population biology of crucifer feeding flea beetles (Coleoptera: Chrysomelidae). Ph.D. Thesis, Dep. of Entomol. Cornell Univ., 106 p.

Tahvanainen, J.O. 1972. Phenology and microhabitat selection of some flea beetles (Coleoptera: Chrysomelidae) on wild and cultivated crucifers in Central New York. Entomol. Scand. 3: 120-38.

Tahvanainen, J.O. 1983. The relationship between flea beetles and their cruciferous host plants: The role of plant and habitat characteristics. Oikos 40: 433-37. (In English; Russian summary).

Tahvanainen, J.O. and R.B. Root. 1970. The invasion and population outbreak of Psylliodes napi (Coleoptera: Chrysomelidae) on yellow rocket (Barbarea vulgaris) in New York. Ann. Entomol. Soc. Am. 63: 1479-80.

Tahvanainen, J.O. and R.B. Root. 1972. The influence of vegetational diversity on the population ecology of a specialized herbivore, Phyllotreta cruciferae (Coleoptera: Chrysomelidae). Oecologia (Berl.) 10: 321-46.

Tandon, H.L.S. 1973. Crop nutrition-pest incidence complex in India. PANS 19: 372-80.

Taylor, T.H. 1918. . Observations on the habits of the turnip flea-beetle. Entomologist (Lond.) 1918: 83-86.

Taylor, W.E. and R. Bardner. 1968. Effects of feeding by larvae of Phaedon cohleanae (F.) and Plutella maculipennis on the yield of radish and turnip plants. Ann. Appl. Biol. 672: 249-54.

Thakur, D.K. 1976. Pest complex of mustard crop and relative efficacy of some important insecticides against important insect pests of mustard crop. Thesis Abstr. Haryana Agric. Univ. 2(2): 119.

Thompson, A.R., R.B. Maude, A.L. Percivall and J.M.. Bambridge. 1982. Evaluation of insecticides for protection against flea beetles of cabbage seedlings raised from seed treated with the fungicide iprodione. Ann. Appl. Biol. 100 (Suppl.): 16-17.

Thorsteinson, A.J. 1960. Infestation flights of sweet clover weevils and flea beetles. Manit. Agron. Annu. Conf., 1960: 31.

Tiittanen, K. and A-L. Varis. 1960. The treatment of seeds of swede, turnip, and turnip rape in the control of flea beetles (Phyllotreta spp.) and cabbage root flies (Hylemia brassicae Bouché and H. floralis Fall.). Publ. Finnish State Agric. Res. Board No. 181: 1-11. (Finnish summary).

Tiittanen, K. and A-L. Varis. 1963. The effect of storage on the germination of lindane-treated seeds and on the efficacy of such treatment in controlling flea beetles (Phyllotreta spp.) and cabbage root flies (Hylemia spp.). Ann. Agric. Fenn. 2: 44-45.

Tokunaga, M. and S. Kadowaki. 1949. Studies on the life history and bionomics of Phyllotreta vittata Fabricius (Col. Chrysomelidae). Feeding habits of imaginal insects. Trans. Kansai Entomol. Soc. 14(2): 59-69.

Tokunaga, M. and S. Kadowaki. 1950a. Studies on the life history and bionomics of Phyllotreta vittata Fabricius. Feeding habits of imaginal insects. Food plants of common Phyllotreta species. Trans. Kansai Entomol. Soc. 15(1): 63-72.

Tokunaga, M. and S. Kadowaki. 1950b. Studies on the life history and bionomics of Phyllotreta vittata Fabricius: Feeding habits of imaginal insects (continued). Trans. Kansai Entomol. Soc. 15(2): 61-68.

Tolg, F. 1938. Determination of the host plants of small Asiatic Halticinae and Ceuthorrhynchinae. Kolon. Rundsch. 24: 234-41. (In German).

Turnock, W.J. and R.J. Lamb. 1982. Protecting canola from flea beetle damage. Can. Agric. 27(4): 4-9. (In English; French).

Twinn, C.R. 1945. A summary of the more important insect conditions in Canada in 1945. Rep. Entomol. Soc. Ont., 76, Ottawa, 1945: 49-55.

Vappula, N.A. 1962. Pests of cultivated plants in Finland. Ann. Agric. Fenn. Vol. 1. Anim. Nocentia No. 5 (suppl. 1): 1-239.

Vargas, P. and W.J.S. Kershaw. 1979. Host selection and choice of feeding site by the flea beetle Phyllotreta undulata Kutsch. An. Inst. Nac. Invest. Agrar. ser. Prot. Veg. (10): 81-91. (In English; Spanish).

Varis, A-L. 1958. On the importance of the sowing time of crucifers in preventing the damage caused by cabbage maggots (Hylemia spp.), on the basis of tests made on big-leafed turnip. Maaloustiet. Aikak. 30: 264-70.

Varis, A-L. and K. Tiittanen. 1982. Insect pests of vegetables in Finland and methods for their control. Acta Entomol. Fenn. 40(0): 38-42.

Varma, B.K. 1961. Bionomics of Phyllotreta cruciferae Goeze (Coleoptera; Chrysomelidae). Indian. J. Agric. Sci. 31: 59-63.

Vincent, C. 1982. Bionomics and field behavior of crucifer-feeding flea beetle adults (Coleoptera: Chrysomelidae). Ph.D. Thesis, Dept. of Entomology, McGill Univ., 172 p.

Vincent, C. and R.K. Stewart. 1981. Altises (Coleoptera: Chrysomelidae) associées aux crucifères cultivées dans le Sud-Ouest du Québec. Ann. Soc. Entomol. Qué. 26: 112-18.

Vincent, C. and R.K. Stewart. 1983. Crucifer-feeding flea beetle dispersal and statistics of directional data. Environ. Entomol. 12: 1380-83.

Vincent, C. and R.K. Stewart. 1984. Effect of allyl isothiocyanate on field behavior of crucifer-feeding flea beetles (Coleoptera: Chrysomelidae). J. Chem. Ecol. 10: 33-39.

Walsh, B.D. and C.V. Riley. 1869. The wavy-striped flea-beetle (Haltica (Phyllotreta) striolata, Illiger). Am. Entomol. 1: 158-59.

Walton, C.L. 1935. The control of flea beetles by means of a seed dressing. Rep. Long Ashton Agric. Hort. Res. Stn. Univ. Bristol. 1935: 80-86.

Warchalowski, A. 1973. Second contribution to the knowledge of the Halticinae of Northern Persia (Coleoptera: Chrysomelidae). Pol. Pismo Entomol. 43: 659-87. (In German).

Warchalowski, A. 1974a. Halticinae (Coleoptera: Chrysomelidae) captured in Mongolia. Fragm. Faun. (Warsaw) 20: 53-60. (In German).

Warchalowski, A. 1974b. Survey of the leaf beetles of Bulgaria (Coleoptera: Chrysomelidae). Pol. Pismo Entomol. 44: 473-542. (In German).

Weires, R.W. and H.C. Chiang. 1973. Integrated control prospects of major cabbage insect pest in Minnesota - based on the faunistic, host

varietal, and trophic relationships. Univ. Minn. Agric. Exp. Stn. Tech. Bull. No. 291, 42 p.

Westdal, P.H. 1950. The life history and behavior of the flea beetle Phyllotreta pusilla Horn in Manitoba. M.Sc. Thesis, Dep. of Zoology, Univ. of Manitoba, Winnipeg, 55 p.

Westdal, P.H. and W. Romanow. 1972. Observations on the biology of the flea beetle, Phyllotreta cruciferae (Coleoptera: Chrysomelidae). Manit. Entomol. 6: 35-45.

White, R.E. 1970. Authorship of the genus Phyllotreta (Chrysomelidae: Coleoptera). Proc. Entomol. Soc. Wash. 72: 388-89.

Wilcox, J.A. 1954. Leaf beetles of Ohio (Chrysomelidae: Coleoptera). Ohio State Univ. Stud., Ohio Biol. Surv. Bull. 43 (Vol. 8, No. 3): 353-506.

Wilcox, J.A. 1979. Leaf beetle host plants in northeastern North America. (Coleoptera: Chrysomelidae). Publ. for Biol. Res. Inst. North Am. by World Natural History Publications, Kinderhook, N.Y.. 30 p.

Winn, A.F. 1910. The horse-radish flea-beetle (Phyllotreta armoraciae, ____ Koch). Annu. Rep. Entomol. Soc. Ont. 41: 59-60.

Wylie, H.G. 1979. Observations on distribution, seasonal life history, and abundance of flea beetles (Coleoptera: Chrysomelidae) that infest rape crops in Manitoba. Can. Entomol. 111: 1345-53.

Wylie, H.G. 1980. Factors affecting facultative diapause of Microctonus vittatae (Hymenoptera: Braconidae). Can. Entomol. 112: 747-49.

Wylie, H.G. 1981. Effects of collection method on estimates of parasitism and sex ratio of flea beetles (Coleoptera: Chrysomelidae) that infest rape crops in Manitoba. Can. Entomol. 113: 665-71.

Wylie, H.G. 1982. An effect of parasitism by Microctonus vittatae (Hymenoptera: Braconidae) on emergence of Phyllotreta cruciferae and Phyllotreta striolata (Coleoptera: Chrysomelidae) from overwintering sites. Can. Entomol. 114: 727-32.

Wylie, H.G. 1983a. Delayed development of Microctonus vittatae (Hymenoptera: Braconidae) in superparasitized adults of Phyllotreta cruciferae (Coleoptera: Chrysomelidae). Can. Entomol. 115: 441-42.

Wylie, H.G. 1983b. Oviposition and survival of the European parasite Microctonus bicolor (Hymenoptera: Braconidae) in crucifer-infesting flea beetles (Coleoptera: Chrysomelidae) in Manitoba. Can. Entomol. 115: 55-58.

Wylie, H.G. 1984. Oviposition and survival of three Nearctic euphorine braconids in crucifer-infesting flea beetles (Coleoptera: Chrysomelidae). Can. Entomol. 116: 1-4.

Wylie, H.G. and C. Loan. 1984. Five Nearctic and one introduced euphorine species that parasitize adults of crucifer-infesting flea beetles (Coleoptera: Chrysomelidae). Can. Entomol. 116: 235-46.

Wylie, H.G., W.J. Turnock and L. Burgess. 1984. Phyllotreta spp., flea beetles (Coleoptera: Chrysomelidae), p. 73-76. In Kelleher, J.S. and M.A. Hulme, (eds), Biological control programs against insects and weeds in Canada 1968-1980. Tech. Commun. Commonwealth Inst. Biol. Control No. 8. 410 p.